

Contents lists available at [SciVerse ScienceDirect](http://www.sciencedirect.com)

Combinatorial Chemistry - An Online Journal

journal homepage: www.elsevier.com/locate/comche

Combinatorial Chemistry Online

Volume 14, Issue 2, February 2012

N.K. Terrett

Ensemble Therapeutics Corp., Cambridge, MA 02139, USA

1. Current literature highlights


1.1. The cascade reaction of isatins with heterocyclic ketene amins

Quinoline derivatives often have biological or pharmacological activity and thus are important molecules in drug discovery. Over the years, many methods have been developed to permit synthetic access, and the importance of these routes has been confirmed by the regular use of named reactions such as the Skraup, Doebner-von Miller, Friedländer, Pfitzinger, Conrad–Limpach and Combes syntheses. The biological and pharmacological properties of quinolines include anti-cancer, antituberculosis, antifungal, and anti-inflammatory activities, and many other compounds act as ligands for adenosine receptors, agonists of P-selectin and inhibitors of caspases.

However, the preparative approaches to these molecules are frequently multi-step, and there remains a need for concise and efficient approaches to quinoline synthesis. There has been a recent rapid growth in interest in one pot transformations, wherein complex molecular scaffolds can be readily built up through cascade chemistry, saving considerable synthetic time and effort. A recent publication describes the reaction of heterocyclic ketene acetals (HKAs) with isatins to generate novel imidazopyrroloquinoline derivatives in good to excellent yield.¹

Within the HKAs (**2**), the conjugation of electron-donating amino groups and the electron-withdrawing carbonyl through a polarised double bond, results in high electron density on the α -carbon and the two secondary amino groups (N1 and N5). Thus they can act as bis-nucleophiles and react rapidly with bis-electrophiles such as isatins (**1**) to generate fused heterocyclic products (**3**).

The reaction to generate imidazopyrroloquinoline derivatives from 5-membered HKAs (**2**, $n = 1$) was investigated in a range of solvents employing various catalysts including zinc chloride and



trifluoroacetic acid. Ultimately it was found that the best yields were obtained using acetic acid in toluene under reflux for 4 h. A number of other five-membered HKAs were used as substrates to react with isatins, and then the reaction was also extended to six- (**2**, $n = 2$) and seven-membered (**2**, $n = 3$) HKAs, building a library of 27 variants. Overall, the compounds were produced in good to excellent yields with very high regioselectivity, using an efficient cascade reaction highly suitable for combinatorial synthesis.

E-mail: nterrett@ensemlediscovery.com

2. A summary of the papers in this month's issue

2.1. Solid-phase synthesis

A multistep approach to construct novel 3-(1*H*-benzo[d]imidazol-2-yl)imidazolidine-2,4-diones and 3-(1*H*-benzo[d]imidazol-2-yl)-2-thioxoimidazolidin-4-ones from commercially available amino acids, amines, and carboxylic acids has been described. The desired aminobenzimidazole tethered hydantoins or thiohydantoins were isolated in good yields.²

An efficient method for the solid-phase synthesis of hydroxamic acids has been described. The method comprises the nucleophilic displacement of esters immobilized on PEGA resins with hydroxylamine/sodium hydroxide in isopropanol. The hydroxylamineolysis protocol is compatible with a broad range of PEGA-supported peptide and peptidomimetic esters, and also found to be compatible with two new strategies for the synthesis of solid-supported lactams and diketopiperazines.³

A recent paper describes a convenient method for the direct amidation of methyl and ethyl β -ketoesters to generate solid-supported β -ketohydroxamates or β -ketoamides and the application of this methodology to the synthesis of 1,5-disubstituted pyrazole-4-hydroxamic acids and pyrazole-4-carboxamides.⁴

2.2. Solution-phase synthesis

A novel multicomponent assembly process-cycloaddition sequence has been applied to the facile synthesis of β -carboline intermediates to gain rapid access to novel derivatives of Yohimbine-like and Corynanthe-like compounds. These products may be easily diversified by cross-coupling reactions and *N*-derivatisations to generate small compound libraries.⁵

2.3. Scaffolds and synthons for combinatorial libraries

No papers this month.

2.4. Solid-phase supported reagents

The design and synthesis of a novel and selective SH-group biotinylating reagent, KSH-1, for the biotinylation of small molecules using solid-phase chemistry, has been described. The results demonstrate that this reagent efficiently biotinylated the small molecule, captopril, and afforded products in high yield and purity.⁶

Silica-supported boron trifluoride (BF₃·SiO₂) has been used as an efficient and reusable catalyst for the synthesis of highly diversified tetrahydropyridines by one-pot multicomponent reactions of amines and aldehydes with β -ketoester. This method proceeds through the intermolecular Mannich reaction followed by intramolecular Mannich reaction. This method has the advantage of easy work-up, convenient, relatively short reaction times and the products were isolated with high yields.⁷

2.5. Novel resins, linkers and techniques

Fragment-based lead discovery is a drug discovery approach that has emerged in the past decade. Because the initial fragments identified in screening typically show weak binding affinity, an intensive medicinal chemistry effort would be required to grow initial fragments into a potential lead compound. A recent publication demonstrates a kinase focused evolved fragment (KFEF) library, constructed by click chemistry-based fragment assembly, that is a valuable source of kinase inhibitors. The screening of this triazole-based KFEF library allowed the rapid identification of potent lead candidates for FLT3 and GSK3 β kinase.⁸

2.6. Library applications

Libraries of dibasic compounds designed around the molecular scaffold of the DA₂/ β_2 dual agonist sibenadet (Viozan™) have yielded a number of promising starting points that have been further optimised into novel potent and selective target molecules with required pharmacokinetic properties. One compound was selected as a novel, high potency, and highly efficacious β_2 -agonist with high selectivity and a duration of action commensurable with once daily dosing.⁹

A novel series of pyrrolidine derived BACE-1 inhibitors have been developed. The potency of the weak initial lead structure was enhanced using library-based SAR methods, and the series was further advanced by rational design while maintaining a minimal ligand binding efficiency threshold. Ultimately, the co-crystal structure was obtained revealing that these inhibitors interacted with the enzyme in a unique fashion and allowed for potent binding in a nontraditional fashion.¹⁰

A recent study has achieved the design and diversity-oriented synthesis of novel 1,4-thiazepine derivatives. These compounds are embedded with a carbazole, pyrazole or isoxazole motif via microwave-assisted multicomponent reactions under solvent-free condition. The compounds have been subjected to testing for *in vitro* antioxidant and cytotoxic activities, resulting in the finding that 1,4-thiazepine derivatives not only have significant antioxidant activity, but also exhibit remarkably selective cytotoxicity to carcinoma cell line HCT 116.¹¹

Thirty-one 2'-hydroxychalcones have been prepared via solid-phase synthesis using base-catalysed aldol condensation of substituted 2'-hydroxyacetophenones and benzaldehydes. Chalcones were tested for their growth inhibitory activity in three human tumour cell lines (MCF-7, NCI-H460 and A375-C5) and results revealed that several of the tested compounds caused a pronounced dose-dependent growth inhibitory effect on the tumour cell lines.¹²

Naturally occurring peptides often possess a macrocyclic and *N*-methylated backbone. These features grant them structural rigidity, high affinity to targets, proteolytic resistance, and occasionally membrane permeability. Because such peptides are produced by either nonribosomal peptide synthetases or enzymatic post-translational modifications, it is a formidable challenge to prepare libraries to screen for bioactivity. A new means of synthesising a *de novo* library of "natural product-like" macrocyclic *N*-methyl-peptides has been described. The selection of anti-E6AP macrocyclic *N*-methyl-peptides, has resulted in the identification of one compound which strongly inhibits polyubiquitination of proteins such as p53.¹³

References

1. Yu F, Yan S, Hu L, Wang Y, Lin J. Cascade reaction of isatins with heterocyclic ketene aminals: synthesis of imidazopyrroloquinoline derivatives. *Org Lett* 2011;**13**(18):4782–5.
2. Dadiboyena S, Nefzi A. Parallel solid-phase synthesis of disubstituted 3-(1*H*-benzodimidazol-2-yl)imidazolidine-2, 4-diones and 3-(1*H*-benzodimidazol-2-yl)-2-thioxoimidazolidin-4-ones. *Tetrahedron Lett* 2011;**52**(52):7030–3.
3. Nandurkar NS, Petersen R, Qvortrup K, Komnatnyy VV, Taveras KM, Le Quement ST, et al. A convenient procedure for the solid phase synthesis of hydroxamic acids on PEGA resins. *Tetrahedron Lett* 2011;**52**(52):7121–4.
4. Zhai W, Gerritz SW, Sofia MJ. Solid phase synthesis of 1, 5-disubstituted pyrazole-4-hydroxamic acids and pyrazole-4-carboxamides via direct amidation of β -ketoesters. *Tetrahedron Lett* 2012;**53**(3):267–70.
5. Wang Z, Kaneda K, Fang Z, Martin SF. Diversity oriented synthesis: concise entry to novel derivatives of Yohimbine and Corynanthe alkaloids. *Tetrahedron Lett* 2012;**53**(5):477–9.
6. Fukumoto K, Adachi K, Kajiyama A, Yamazaki Y, Yakushiji F, Hayashi Y. Development of a solid-supported biotinylation reagent for efficient biotin labeling of SH groups on small molecules. *Tetrahedron Lett* 2012;**53**(5):535–8.
7. Ramachandran R, Jayanthi S, Jeong YT. One-pot synthesis of highly diversified tetrahydropyridines by tandem condensation of aldehydes, amines, and β -ketoesters. *Tetrahedron* 2012;**68**(1):363–9.

8. Irie T, Fujii I, Sawa M. Design and combinatorial synthesis of a novel kinase-focused library using click chemistry-based fragment assembly. *Bioorg Med Chem Lett* 2012;**22**(1):591–6.
9. Alcaraz L, Bailey A, Cadogan E, Connolly S, Jewell R, Jordan S, et al. From libraries to candidate: the discovery of new ultra long-acting dibasic β_2 -adrenoceptor agonists. *Bioorg Med Chem Lett* 2012;**22**(1):689–95.
10. Stachel SJ, Steele TG, Petrocchi A, Haugabook SJ, McGaughey G, Holloway MK, et al. Discovery of pyrrolidine-based β -secretase inhibitors: lead advancement through conformational design for maintenance of ligand binding efficiency. *Bioorg Med Chem Lett* 2012;**22**(1):240–4.
11. Shi F, Zeng X-N, Cao X-D, Zhang S, Jiang B, Zheng W-F, et al. Design and diversity-oriented synthesis of novel 1, 4-thiazepan-3-ones fused with bioactive heterocyclic skeletons and evaluation of their antioxidant and cytotoxic activities. *Bioorg Med Chem Lett* 2012;**22**(1):743–6.
12. Neves MP, Cravo S, Lima RT, Vasconcelos MH, Nascimento MSJ, Silva AMS, et al. Solid-phase synthesis of 2'-hydroxychalcones. Effects on cell growth inhibition, cell cycle and apoptosis of human tumor cell lines. *Bioorg Med Chem* 2012;**20**(1):25–33.
13. Yamagishi Y, Shoji I, Miyagawa S, Kawakami T, Katoh T, Goto Y, et al. Natural product-like macrocyclic N-methyl-peptide inhibitors against a ubiquitin ligase uncovered from a ribosome-expressed de novo library. *Chem Biol* 2011;**18**(12):1562–70.
- Leung D, Kang SO, Anslyn EV. Rapid determination of enantiomeric excess: a focus on optical approaches. *Chem Soc Rev* 2012;**41**(1):448–79.
- Guido RVC, Oliva G, Andricopulo AD. Modern drug discovery technologies: opportunities and challenges in lead discovery. *Comb Chem High Throughput Screening* 2011;**14**(10):830–9.
- Barua S, Ramos J, Potta T, Taylor D, Huang H-C, Montanez G, et al. Discovery of cationic polymers for non-viral gene delivery using combinatorial approaches. *Comb Chem High Throughput Screening* 2011;**14**(10):908–24.
- McGee JE, Roy A. Patent review. *Comb Chem High Throughput Screening* 2011;**14**(10):926–8.
- Lau WF, Withka JM, Hepworth D, Magee TV, Du YJ, Bakken GA, et al. Design of a multi-purpose fragment screening library using molecular complexity and orthogonal diversity metrics. *J Comput Aided Mol Des* 2011;**25**(7):621–36.
- Xiang J, Geng C, Yi L, Dang Q, Bai X. Synthesis of highly substituted 2, 3-dihydropyrimido[4,5-d]pyrimidin-4(1H)-ones from 4, 6-dichloro-5-formylpyrimidine, amines and aldehydes. *Mol Diversity* 2011;**15**(4):839–47.
- Kouznetsov VV, Gomez CMM, Parada LKL, Bermudez JH, Mendez LYV, Acevedo AM. Efficient synthesis and free-radical scavenging capacity of new 2,4-substituted tetrahydroquinolines prepared via BiCl₃-catalyzed three-component Povarov reaction, using N-vinylamides. *Mol Diversity* 2011;**15**(4):1007–16.
- Mabkhot YN, Al-Majid AM, Barakat A, Alshahrani S, Siddiqui Y. 1,1'-(3-Methyl-4-phenylthieno[2,3-b]thiophene-2,5-diyl)diethanone as a building block in heterocyclic synthesis. Novel synthesis of some pyrazole and pyrimidine derivatives. *Molecules (Basel, Switzerland)* 2011;**16**(8):6502–11.
- Basso A, Banfi L, Riva R. Divergent synthesis of novel five-membered heterocyclic compounds by base-mediated rearrangement of acrylamides derived from a novel isocyanide-based multicomponent reaction. *Molecules (Basel, Switzerland)* 2011;**16**(10):8775–87.
- Thompson CM, Poole JL, Cross JL, Akritopoulou-Zanze I, Djuric SW. Small molecule library synthesis using segmented flow. *Molecules (Basel, Switzerland)* 2011;**16**(11):9161–77.
- White TR, Renzelman CM, Rand AC, Rezai T, McEwen CM, Gelev VM, et al. On-resin N-methylation of cyclic peptides for discovery of orally bioavailable scaolds. *Nat Chem Biol* 2011;**7**(11):810–7.
- Miljanic OS. Reversible covalent chemistry: alcohols in the mix. *Nat Chem* 2011;**3**(12):909–10.
- Montavon TJ, Li J, Cabrera-Pardo JR, Mrksich M, Kozmin SA. Three-component reaction discovery enabled by mass spectrometry of self-assembled monolayers. *Nat Chem* 2012;**4**(1):45–51.
- Hardin Narayan AR, Sarpong R. Indolizinones as synthetic scaolds: fundamental reactivity and the relay of stereochemical information. *Org Biomol Chem* 2012;**10**(1):70–8.
- Estrada AA, Lyssikatos JP, St-Jean F, Bergeron P. Access to saturated fused pyrimidine derivatives via a flexible N-vinyl tertiary enamide synthesis. *Synlett* (16):2387–91.

Further reading

Papers on combinatorial chemistry or solid-phase synthesis from other journals

- Dobele M, Wiehn MS, Brase S. Traceless solid-phase synthesis of trifluoromethylarenes. *Angew Chem (International ed. in English)* 2011;**50**(48):11533–5.
- Kim C-B, Cho C-H, Jo MJ, Park K. Solid-phase synthesis of unfunctionalized arenes via the traceless cleavage of sulfonate linkers. *Bull Korean Chem Soc* 2011;**32**(10):3655–9.
- Takayama H, Takahashi S, Moriya T, Osada H, Iwabuchi Y, Kanoh N. Detection of cytochrome P450 substrates by using a small-molecule droplet array on an NADH-immobilized solid surface. *ChemBioChem* 2011;**12**(18):2748–52.
- Bartoloni M, Kadam RU, Schwartz J, Furrer J, Darbre T, Reymond J-L. Expanding the accessible chemical space by solid phase synthesis of bicyclic homodetic peptides. *Chem Commun (Cambridge, United Kingdom)* 2011;**47**(47):12634–6.
- Mannocci L, Leimbacher M, Wichert M, Scheuermann J, Neri D. 20 years of DNA-encoded chemical libraries. *Chem Commun (Cambridge, United Kingdom)* 2012;**47**(48):12747–53.
- Daguer JP, Ciobanu M, Alvarez S, Barluenga S, Winssinger N. DNA-templated combinatorial assembly of small molecule fragments amenable to selection/amplification cycles. *Chem Sci* 2011;**2**(4):625–32.